

- [4] Blennerhassett JM, Jayalath VM. The Four Square Step Test is a feasible and valid clinical test of dynamic standing balance for use in ambulant people poststroke[J]. Arch Phys Med Rehabil, 2008, 89(11):2156—2161.
- [5] Duncan RP, Earhart GM. Four square step test performance in people with Parkinson disease[J]. J Neurol Phys Ther, 2013, 37(1):2—8.
- [6] Wagner JM, Norris RA, Van Dillen LR, et al. Four Square Step Test in ambulant persons with multiple sclerosis: validity, reliability, and responsiveness[J]. Int J Rehabil Res, 2013, 36(3):253—259.
- [7] Bandong AN, Madriaga GO, Gorgon EJ. Reliability and validity of the Four Square Step Test in children with cerebral palsy and Down syndrome[J]. Res Dev Disabil, 2015, (47):39—47.
- [8] Dite W, Connor HJ, Curtis HC. Clinical identification of multiple fall risk early after unilateral transtibial amputation [J]. Arch Phys Med Rehabil, 2007, 88(1):109—114.
- [9] Said CM, Goldie PA, Patla AE, et al. Obstacle crossing in subjects with stroke[J]. Arch Phys Med Rehabil, 1999, 80(9): 1054—1059.
- [10] Hyndman D, Ashburn A, Stack E. Fall events among people with stroke living in the community: circumstances of falls and characteristics of fallers[J]. Arch Phys Med Rehabil, 2002, 83(2):165—170.
- [11] Blennerhassett JM, Dite W, Ramage ER, et al. Changes in balance and walking from stroke rehabilitation to the community: a follow-up observational study[J]. Arch Phys Med Rehabil, 2012, 93(10):1782—1787.
- [12] Goh EY, Chua SY, Hong SJ, et al. Reliability and concurrent validity of Four Square Step Test scores in subjects with chronic stroke: a pilot study[J]. Arch Phys Med Rehabil, 2013, 94(7):1306—1311.
- [13] Speelman AD, van de Warrenburg BP, van Nimwegen M, et al. How might physical activity benefit patients with Parkinson disease?[J]. Nat Rev Neurol, 2011, 7(9):528—534.
- [14] Paul SS, Sherrington C, Fung VS, et al. Motor and cognitive impairments in Parkinson disease: relationships with specific balance and mobility tasks[J]. Neurorehabil Neural Repair, 2013, 27(1):63—71.
- [15] McKee KE, Hackney ME. The Four Square Step Test in individuals with Parkinson's disease: association with executive function and comparison with older adults[J]. Neuro Rehabilitation, 2014, 35(2):279—289.
- [16] Nilsagard YE, von Koch LK, Nilsson M, et al. Balance exercise program reduced falls in people with multiple sclerosis: a single-group, pretest-posttest trial[J]. Arch Phys Med Rehabil, 2014, 95(12):2428—2434.
- [17] Ilett P, Lythgo N, Martin C, et al. Balance and gait in people with multiple sclerosis: A comparison with healthy controls and the immediate change after an intervention based on the Bobath concept[J]. Physiother Res Int, 2015, [Epub ahead of print].
- [18] Kloos AD, Fritz NE, Kostyk SK, et al. Clinimetric properties of the Tinetti Mobility Test, Four Square Step Test, Activities-specific Balance Confidence Scale, and spatiotemporal gait measures in individuals with Huntington's disease [J]. Gait Posture, 2014, 40(4):647—651.
- [19] Gorgon EJ, Madriaga GO, Gomez-Cailao MZ, et al. Reliability and feasibility of the four square step test for use in children with cerebral palsy: a pilot study[J]. The Internet Journal of Allied Health Sciences and Practice, 2014, 12(2):1—5.
- [20] Verma A, Samuel AJ, Aranha VP. The four square step test in children with down syndrome: Reliability and concurrent validity[J]. J Pediatr Neurosci, 2014, 9(3):221—226.
- [21] MacDuff GS, Krantz PJ, McClannahan LE. Prompts and prompt-fading strategies for people with autism. In: Making a difference. edn. Edited by C. Maurice, G. Green, Foxx RM. Austin: TX: Pro-Ed, 2001: 37—50.

北京大学第一届全国运动神经元病诊治技术新进展学习班招生通知

北京大学第三医院神经内科主办的“北京大学第一届全国运动神经元病(MND)诊治技术新进展学习班”拟于2017年7月17—21日在北京举行。本学习班集中了北京大学神经病学诊治领域雄厚的师资力量,并邀请国内知名专家共同参与授课。学习班内容包括MND的基础研究,临床电生理技术、认知筛查等在MND的诊断治疗中的应用,以及MND相关的营养支持、呼吸管理、精神心理支持及康复等,欢迎全国神经内科、电生理检查室、康复科、精神心理科及其他相关科室的医技术人员参加。国家级继续医学教育I类8学分。

E-mail: bssn1706@sina.com, 联系电话: 15901312366, 张华纲; 01082264446, 张朔。

北京大学第三医院神经内科